

Yarr., is considered identical with *Polyprion cernuum*, Cuv. and Vall. *Hemulon formosum* or squirrel fish, a native of the West Indian Seas, has been once taken. *Sciæna Aquila* has been taken twice. *Cyclopterus coronatus*, coronated Lump fish of Couch, and considered new to science—one specimen however has only been examined, of a very small size; see our Miscellanea for characters. At the conclusion of this list there are some important remarks on the time and œconomy of spawning of many of the fishes commercially used, together with some hints regarding the improvement of our fishing regulations. This is a much more important subject than most people are at present willing to suppose, and these remarks deserve attention.

Of the Crustaceous animals, fifty-four species are enumerated; five *Lepadæa*, and of the Radiated animals, twenty-three. We trust the second part of this Fauna will speedily appear.

British Entomology. By J. Curtis, F.L.S.

Since we last noticed this work Nos. 175 to 180 have been published, which complete the 15th volume.

Lampyrus noctiluca (the Glow-worm), *Rhagio Heyshami*, *Ephemerella cognata* (the May-fly), *Harpocera Burmeisteri*, *Libellula rubicunda*, *Anthicus tibialis*, *Molanna angustata*, *Coccus Aceris*, and *Phytosus spinifer* are amongst the most interesting novelties, and there is a very beautiful plate of the rare *Clostera anachoreta* and its caterpillar. We would also call the attention of botanists to the faithful figures of *Astragalus hypoglottis*, *Sagittaria sagittifolia*, *Arabis stricta*, *Osmunda regalis*, *Althæa officinalis*, *Milium effusum*, *Hutchinsia petraea*, *Oxyria reniformis*, *Acorus Calamus*, *Zannichellia palustris*, *Melittis grandiflora*, &c.

We are happy to find that Mr. Curtis is preparing for the press a SYNOPSIS OF BRITISH INSECTS, the orders to be published in separate volumes. This will undoubtedly be an agreeable present to all lovers of entomology, especially those who have not ready access to large libraries.

Entomologists are earnestly invited to supply Mr. Curtis with duplicate specimens of those insects which he does not possess, or to lend him such species as maybe required for describing.

Icones Plantarum. By Sir W. J. Hooker.

The fifth part of this work, or the first half of the third volume, has recently appeared, with fifty plates, from Tabs. CCI. to CCL. inclusive. This portion is particularly rich in new South American plants from the collections of Mr. Gardner, Mathews, Professor W. Jameson, &c.

Yarr., is considered identical with *Polyprion cernuum*, Cuv. and Vall. *Hemulon formosum* or squirrel fish, a native of the West Indian Seas, has been once taken. *Sciæna Aquila* has been taken twice. *Cyclopterus coronatus*, coronated Lump fish of Couch, and considered new to science—one specimen however has only been examined, of a very small size; see our Miscellanea for characters. At the conclusion of this list there are some important remarks on the time and œconomy of spawning of many of the fishes commercially used, together with some hints regarding the improvement of our fishing regulations. This is a much more important subject than most people are at present willing to suppose, and these remarks deserve attention.

Of the Crustaceous animals, fifty-four species are enumerated; five *Lepadæa*, and of the Radiated animals, twenty-three. We trust the second part of this Fauna will speedily appear.

British Entomology. By J. Curtis, F.L.S.

Since we last noticed this work Nos. 175 to 180 have been published, which complete the 15th volume.

Lampyris noctiluca (the Glow-worm), *Rhagio Heyshami*, *Ephemeræ cognata* (the May-fly), *Harpocera Burmeisteri*, *Libellula rubicunda*, *Anthicus tibialis*, *Molanna angustata*, *Coccus Aceris*, and *Phytosus spinifer* are amongst the most interesting novelties, and there is a very beautiful plate of the rare *Clostera anachoreta* and its caterpillar. We would also call the attention of botanists to the faithful figures of *Astragalus hypoglottis*, *Sagittaria sagittifolia*, *Arabis stricta*, *Osmunda regalis*, *Althæa officinalis*, *Milium effusum*, *Hutchinsia petraea*, *Oxyria reniformis*, *Acorus Calamus*, *Zannichellia palustris*, *Melittis grandiflora*, &c.

We are happy to find that Mr. Curtis is preparing for the press a SYNOPSIS OF BRITISH INSECTS, the orders to be published in separate volumes. This will undoubtedly be an agreeable present to all lovers of entomology, especially those who have not ready access to large libraries.

Entomologists are earnestly invited to supply Mr. Curtis with duplicate specimens of those insects which he does not possess, or to lend him such species as maybe required for describing.

Icones Plantarum. By Sir W. J. Hooker.

The fifth part of this work, or the first half of the third volume, has recently appeared, with fifty plates, from Tabs. CCI. to CCL. inclusive. This portion is particularly rich in new South American plants from the collections of Mr. Gardner, Mathews, Professor W. Jameson, &c.

Yarr., is considered identical with *Polyprion cernuum*, Cuv. and Vall. *Hemulon formosum* or squirrel fish, a native of the West Indian Seas, has been once taken. *Sciæna Aquila* has been taken twice. *Cyclopterus coronatus*, coronated Lump fish of Couch, and considered new to science—one specimen however has only been examined, of a very small size; see our Miscellanea for characters. At the conclusion of this list there are some important remarks on the time and œconomy of spawning of many of the fishes commercially used, together with some hints regarding the improvement of our fishing regulations. This is a much more important subject than most people are at present willing to suppose, and these remarks deserve attention.

Of the Crustaceous animals, fifty-four species are enumerated; five *Lepadæa*, and of the Radiated animals, twenty-three. We trust the second part of this Fauna will speedily appear.

British Entomology. By J. Curtis, F.L.S.

Since we last noticed this work Nos. 175 to 180 have been published, which complete the 15th volume.

Lampyrus noctiluca (the Glow-worm), *Rhagio Heyshami*, *Ephemeræ cognata* (the May-fly), *Harpocera Burmeisteri*, *Libellula rubicunda*, *Anthicus tibialis*, *Molanna angustata*, *Coccus Aceris*, and *Phytosus spinifer* are amongst the most interesting novelties, and there is a very beautiful plate of the rare *Clostera anachoreta* and its caterpillar. We would also call the attention of botanists to the faithful figures of *Astragalus hypoglottis*, *Sagittaria sagittifolia*, *Arabis stricta*, *Osmunda regalis*, *Althæa officinalis*, *Milium effusum*, *Hutchinsia petraea*, *Oxyria reniformis*, *Acorus Calamus*, *Zannichellia palustris*, *Melittis grandiflora*, &c.

We are happy to find that Mr. Curtis is preparing for the press a SYNOPSIS OF BRITISH INSECTS, the orders to be published in separate volumes. This will undoubtedly be an agreeable present to all lovers of entomology, especially those who have not ready access to large libraries.

Entomologists are earnestly invited to supply Mr. Curtis with duplicate specimens of those insects which he does not possess, or to lend him such species as maybe required for describing.

Icones Plantarum. By Sir W. J. Hooker.

The fifth part of this work, or the first half of the third volume, has recently appeared, with fifty plates, from Tabs. CCI. to CCL. inclusive. This portion is particularly rich in new South American plants from the collections of Mr. Gardner, Mathews, Professor W. Jameson, &c.

Yarr., is considered identical with *Polyprion cernuum*, Cuv. and Vall. *Hemulon formosum* or squirrel fish, a native of the West Indian Seas, has been once taken. *Sciæna Aquila* has been taken twice. *Cyclopterus coronatus*, coronated Lump fish of Couch, and considered new to science—one specimen however has only been examined, of a very small size; see our Miscellanea for characters. At the conclusion of this list there are some important remarks on the time and œconomy of spawning of many of the fishes commercially used, together with some hints regarding the improvement of our fishing regulations. This is a much more important subject than most people are at present willing to suppose, and these remarks deserve attention.

Of the Crustaceous animals, fifty-four species are enumerated; five *Lepadæa*, and of the Radiated animals, twenty-three. We trust the second part of this Fauna will speedily appear.

British Entomology. By J. Curtis, F.L.S.

Since we last noticed this work Nos. 175 to 180 have been published, which complete the 15th volume.

Lampyrus noctiluca (the Glow-worm), *Rhagio Heyshami*, *Ephemerella cognata* (the May-fly), *Harpocera Burmeisteri*, *Libellula rubicunda*, *Anthicus tibialis*, *Molanna angustata*, *Coccus Aceris*, and *Phytosus spinifer* are amongst the most interesting novelties, and there is a very beautiful plate of the rare *Clostera anachoreta* and its caterpillar. We would also call the attention of botanists to the faithful figures of *Astragalus hypoglottis*, *Sagittaria sagittifolia*, *Arabis stricta*, *Osmunda regalis*, *Althæa officinalis*, *Milium effusum*, *Hutchinsia petraea*, *Oxyria reniformis*, *Acorus Calamus*, *Zannichellia palustris*, *Melittis grandiflora*, &c.

We are happy to find that Mr. Curtis is preparing for the press a SYNOPSIS OF BRITISH INSECTS, the orders to be published in separate volumes. This will undoubtedly be an agreeable present to all lovers of entomology, especially those who have not ready access to large libraries.

Entomologists are earnestly invited to supply Mr. Curtis with duplicate specimens of those insects which he does not possess, or to lend him such species as maybe required for describing.

Icones Plantarum. By Sir W. J. Hooker.

The fifth part of this work, or the first half of the third volume, has recently appeared, with fifty plates, from Tabs. CCI. to CCL. inclusive. This portion is particularly rich in new South American plants from the collections of Mr. Gardner, Mathews, Professor W. Jameson, &c.

The same author has just completed the second part of Mr. Bauer's 'Illustrations of the Genera of Ferns'; and the seventh part of the 'Botany of Capt. Beechey's Voyage' will soon be ready. These two works, and the 'Flora Boreali-Americana,' of which Part X. is in a state of great forwardness, are published by H. G. Bohn, 4, York Street, Covent Garden.

We have just received the forty-seventh number of Mr. Sowerby's 'Supplement to English Botany.' It contains plates and descriptions of *Polygonum laxum*, Reich. and Borr. in Hook. Brit. Fl., ed. 4. note; *Lotus hispidus*, Loisel, which we have been disposed to consider as not specifically distinct from *L. angustissimus*, and it occurs in Jersey with that species; *Chara pulchella*, Wallr., "principally distinguished from *C. Hedwigii* by its more flexible stems and oblong nucules;" and *Tetraspora lubrica*, Agardh, and Hook. in E. Fl. 5. p. 313.

Tijdschrift voor Natuurlijke Geschiedenis en Physiologie; edited by Prof. J. Van der Hoeven and Prof. W. H. de Vriese, Leiden, 1837.

Part I and II.

These contain the following original articles, besides reviews and notices.

Some remarks on the northern Whale, *Balenoptera rostrata*. By W. Vrolik.—On the Sargasso or Gulf-weed. By F. A. W. Miquel.—Some remarks on the origin of the green colour and changes of form in the stem of plants. By Dr. J. Wttewaall.—Researches respecting the motion of leaves which do not originate from swellings. By M. Dassen.—[The principal results contained in this memoir have been noticed at p. 223. of this Journal.]—Additions to our knowledge of the simple eyes of articulated animals. By A. Brants.—Experiments on the action of poisons on plants. By F. A. W. Miquel.—On the cause of the brand in *Physalia*. By P. W. Korthals.—Some notices of G. R. Treviranus. By J. Van der Hoeven.

Part III. and IV. 1838.

Hints on the origin of monstrous births, and on the doctrine of misformations. By W. Vrolik.—Contributions to the natural history of man. By J. Van der Hoeven.—The vegetation of the Northern Netherlands compared with that of the Prussian Rhine Provinces. By F. A. W. Miquel.—Contributions to the solution of the question, whether *Lemna arrhiza*, auct. be a permanent distinct species, or merely a development form of some other species of the same genus. By J. F. Hoffmann.—On the periodical secretion of blood from the generative organs in some domestic animals, especially in the cow,

The same author has just completed the second part of Mr. Bauer's 'Illustrations of the Genera of Ferns'; and the seventh part of the 'Botany of Capt. Beechey's Voyage' will soon be ready. These two works, and the 'Flora Boreali-Americana,' of which Part X. is in a state of great forwardness, are published by H. G. Bohn, 4, York Street, Covent Garden.

We have just received the forty-seventh number of Mr. Sowerby's 'Supplement to English Botany.' It contains plates and descriptions of *Polygonum laxum*, Reich. and Borr. in Hook. Brit. Fl., ed. 4. note; *Lotus hispidus*, Loisel, which we have been disposed to consider as not specifically distinct from *L. angustissimus*, and it occurs in Jersey with that species; *Chara pulchella*, Wallr., "principally distinguished from *C. Hedwigii* by its more flexible stems and oblong nucules;" and *Tetraspora lubrica*, Agardh, and Hook. in E. Fl. 5. p. 313.

Tijdschrift voor Natuurlijke Geschiedenis en Physiologie; edited by Prof. J. Van der Hoeven and Prof. W. H. de Vriese, Leiden, 1837.

Part I and II.

These contain the following original articles, besides reviews and notices.

Some remarks on the northern Whale, *Balenoptera rostrata*. By W. Vrolik.—On the Sargasso or Gulf-weed. By F. A. W. Miquel.—Some remarks on the origin of the green colour and changes of form in the stem of plants. By Dr. J. Wttewaall.—Researches respecting the motion of leaves which do not originate from swellings. By M. Dassen.—[The principal results contained in this memoir have been noticed at p. 223. of this Journal.]—Additions to our knowledge of the simple eyes of articulated animals. By A. Brants.—Experiments on the action of poisons on plants. By F. A. W. Miquel.—On the cause of the brand in *Physalia*. By P. W. Korthals.—Some notices of G. R. Treviranus. By J. Van der Hoeven.

Part III. and IV. 1838.

Hints on the origin of monstrous births, and on the doctrine of misformations. By W. Vrolik.—Contributions to the natural history of man. By J. Van der Hoeven.—The vegetation of the Northern Netherlands compared with that of the Prussian Rhine Provinces. By F. A. W. Miquel.—Contributions to the solution of the question, whether *Lemna arrhiza*, auct. be a permanent distinct species, or merely a development form of some other species of the same genus. By J. F. Hoffmann.—On the periodical secretion of blood from the generative organs in some domestic animals, especially in the cow,

The same author has just completed the second part of Mr. Bauer's 'Illustrations of the Genera of Ferns'; and the seventh part of the 'Botany of Capt. Beechey's Voyage' will soon be ready. These two works, and the 'Flora Boreali-Americana,' of which Part X. is in a state of great forwardness, are published by H. G. Bohn, 4, York Street, Covent Garden.

We have just received the forty-seventh number of Mr. Sowerby's 'Supplement to English Botany.' It contains plates and descriptions of *Polygonum laxum*, Reich. and Borr. in Hook. Brit. Fl., ed. 4. note; *Lotus hispidus*, Loisel, which we have been disposed to consider as not specifically distinct from *L. angustissimus*, and it occurs in Jersey with that species; *Chara pulchella*, Wallr., "principally distinguished from *C. Hedwigii* by its more flexible stems and oblong nucules;" and *Tetraspora lubrica*, Agardh, and Hook. in E. Fl. 5. p. 313.

Tijdschrift voor Natuurlijke Geschiedenis en Physiologie; edited by Prof. J. Van der Hoeven and Prof. W. H. de Vriese, Leiden, 1837.

Part I and II.

These contain the following original articles, besides reviews and notices.

Some remarks on the northern Whale, *Balenoptera rostrata*. By W. Vrolik.—On the Sargasso or Gulf-weed. By F. A. W. Miquel.—Some remarks on the origin of the green colour and changes of form in the stem of plants. By Dr. J. Wttewaall.—Researches respecting the motion of leaves which do not originate from swellings. By M. Dassen.—[The principal results contained in this memoir have been noticed at p. 223. of this Journal.]—Additions to our knowledge of the simple eyes of articulated animals. By A. Brants.—Experiments on the action of poisons on plants. By F. A. W. Miquel.—On the cause of the brand in *Physalia*. By P. W. Korthals.—Some notices of G. R. Treviranus. By J. Van der Hoeven.

Part III. and IV. 1838.

Hints on the origin of monstrous births, and on the doctrine of misformations. By W. Vrolik.—Contributions to the natural history of man. By J. Van der Hoeven.—The vegetation of the Northern Netherlands compared with that of the Prussian Rhine Provinces. By F. A. W. Miquel.—Contributions to the solution of the question, whether *Lemna arrhiza*, auct. be a permanent distinct species, or merely a development form of some other species of the same genus. By J. F. Hoffmann.—On the periodical secretion of blood from the generative organs in some domestic animals, especially in the cow,

The same author has just completed the second part of Mr. Bauer's 'Illustrations of the Genera of Ferns'; and the seventh part of the 'Botany of Capt. Beechey's Voyage' will soon be ready. These two works, and the 'Flora Boreali-Americana,' of which Part X. is in a state of great forwardness, are published by H. G. Bohn, 4, York Street, Covent Garden.

We have just received the forty-seventh number of Mr. Sowerby's 'Supplement to English Botany.' It contains plates and descriptions of *Polygonum laxum*, Reich. and Borr. in Hook. Brit. Fl., ed. 4. note; *Lotus hispidus*, Loisel, which we have been disposed to consider as not specifically distinct from *L. angustissimus*, and it occurs in Jersey with that species; *Chara pulchella*, Wallr., "principally distinguished from *C. Hedwigii* by its more flexible stems and oblong nucules;" and *Tetraspora lubrica*, Agardh, and Hook. in E. Fl. 5. p. 313.

Tijdschrift voor Natuurlijke Geschiedenis en Physiologie; edited by Prof. J. Van der Hoeven and Prof. W. H. de Vriese, Leiden, 1837.

Part I and II.

These contain the following original articles, besides reviews and notices.

Some remarks on the northern Whale, *Balenoptera rostrata*. By W. Vrolik.—On the Sargasso or Gulf-weed. By F. A. W. Miquel.—Some remarks on the origin of the green colour and changes of form in the stem of plants. By Dr. J. Wttewaall.—Researches respecting the motion of leaves which do not originate from swellings. By M. Dassen.—[The principal results contained in this memoir have been noticed at p. 223. of this Journal.]—Additions to our knowledge of the simple eyes of articulated animals. By A. Brants.—Experiments on the action of poisons on plants. By F. A. W. Miquel.—On the cause of the brand in *Physalia*. By P. W. Korthals.—Some notices of G. R. Treviranus. By J. Van der Hoeven.

Part III. and IV. 1838.

Hints on the origin of monstrous births, and on the doctrine of misformations. By W. Vrolik.—Contributions to the natural history of man. By J. Van der Hoeven.—The vegetation of the Northern Netherlands compared with that of the Prussian Rhine Provinces. By F. A. W. Miquel.—Contributions to the solution of the question, whether *Lemna arrhiza*, auct. be a permanent distinct species, or merely a development form of some other species of the same genus. By J. F. Hoffmann.—On the periodical secretion of blood from the generative organs in some domestic animals, especially in the cow,